

Stellar Envelope Convection calibrated by Radiation Hydrodynamics Simulations: Influence on Globular Cluster Isochrones

Bernd Freytag

Astronomical Observatory, Juliane Maries Vej 30, DK-2100 Copenhagen East, Denmark
[bf@astro.ku.dk],

Institute for Theoretical Physics and Astrophysics, D-24098 Kiel, Germany

and

Maurizio Salaris

Astrophysics Research Institute, Liverpool John Moores University, Twelve Quays House,
Egerton Wharf, Birkenhead L41 1LD, UK [ms@staru1.livjm.ac.uk]

ABSTRACT

One of the largest sources of uncertainty in the computation of globular cluster isochrones and hence in the age determination of globular clusters is the lack of a rigorous description of convection. Therefore, we calibrated the superadiabatic temperature gradient in the envelope of metal-poor low-mass stars according to the results from a new grid of 2D hydrodynamical models, which cover the Main Sequence and the lower Red Giant Branch of globular cluster stars. In practice, we still use for computing the evolutionary stellar models the traditional mixing length formalism, but we fix the mixing length parameter α in order to reproduce the run of the entropy of the deeper adiabatic region of the stellar envelopes with effective temperature and gravity as obtained from the hydro-models. The detailed behaviour of the calibrated α depends in a non-trivial way on the effective temperature, gravity and metallicity of the star. Nevertheless, the resulting isochrones for the relevant age range of galactic globular clusters have only small differences with respect to isochrones computed adopting a constant solar calibrated value of the mixing length. Accordingly, the age of globular clusters is reduced by 0.2 Gyr at most.

Subject headings: convection — stars: evolution — stars: Population II — globular clusters: general

1. Introduction

One of the most important unsolved problems of stellar evolution is the determination of the temperature gradient in the superadiabatic regions at the top of the convective envelopes of cool

stars, which strongly affects the effective temperature (T_{eff}) of these objects. The mixing length theory (MLT - Böhm-Vitense 1958) is widely used for deriving this gradient. It contains a number of free parameters, among them α , the ratio of the mixing length to the pressure scale height, which provides the scale length of the convection. There are different versions of the MLT, each one assuming different values for these parameters. As demonstrated by Pedersen et al. (1990), the T_{eff} values obtained from the different formalisms are equivalent, provided that a suitable value of α is selected (see also Gough & Weiss 1976). This means that the MLT results concerning stellar structure models depend only on one free parameter, namely α , and its absolute value depends on the selected MLT formalism. Once the formalism is fixed, α is usually calibrated by reproducing the solar T_{eff} , and this solar-calibrated α is then used for computing models of stars very different from the Sun (e.g., metal-poor Red Giant Branch and Main Sequence stars of various masses). However, in principle there is no compelling reason that α should be the same for the Sun and different kinds of stars.

More recently, Canuto & Mazzitelli (1991 - CM) proposed a new formalism for the treatment of the superadiabatic convection; they take into account the full turbulent energy spectrum and set the convective scale length equal to the geometrical depth from the top of the convective region. Comparisons between MLT (solar-calibrated α) and CM stellar models show that isochrones computed with the CM formalism cannot be reproduced by the MLT with any constant value of α (Mazzitelli et al. 1995).

The problem of determining accurate effective temperatures for cool stars affects the globular cluster (GC) age determination and, in turn, the estimated age of the universe. Large variations of α alter the derived stellar T_{eff} and colours and change the shape of GC isochrones; as demonstrated by Chaboyer (1995 - see also Chaboyer et al. 1998) the change of the isochrone shape can even modify the luminosity of the Turn Off (TO - bluest point along a given isochrone). Moreover, when comparing CM and MLT isochrones one finds that for the most metal-poor isochrones, in the relevant range of ages of the metal-poor galactic GCs, the TO luminosity obtained from the CM isochrones differs appreciably from the case of the MLT (Mazzitelli et al. 1995). Since the TO brightness is the main age indicator for GCs, uncertainties in the convection treatment can affect the estimated GC ages (by ≈ 1 Gyr or even more).

In principle, one could try to constrain the convective efficiency by comparing theoretical isochrones with regions of the observed Colour-Magnitude diagrams of GCs whose colours are unaffected by the cluster age, but the still existing large uncertainties in the colour-transformations (see, e.g., Weiss & Salaris 1998) do not permit to safely follow this approach. An empirical constraint is given by the T_{eff} of the upper Red Giant Branches of a sample of GCs as derived by Frogel et al. (1981). MLT models computed with a solar calibrated α (see, e.g., Vandenberg et al. 1996, Salaris & Weiss 1998) appear to be in agreement with these empirical data, also if a precise error bar on the α value calibrated in this way is hard to establish, being probably in the range ± 0.1 - 0.3 (see Vandenberg et al. 1996 and references therein). However, this 'empirical' calibration in principle does not constrain the convection along the Main Sequence and the lower

Red Giant Branch of GCs.

Both the MLT and CM formalisms assume a simplified, time-independent, one-dimensional, local treatment of a typically non-stationary, multi-dimensional, and non-local phenomenon which the stellar convection actually is. The final solution to the problem of the superadiabatic convection in stellar envelopes has to come from the computation of realistic multidimensional radiation-hydrodynamics (RHD) simulations covering the range of effective temperatures, gravities, and compositions typical of stars with convective envelopes. First attempts to include in stellar models the results from rather crude 2-dimensional (2D) and 3-dimensional (3D) hydrodynamical simulations date back to the works by Deupree & Varner (1980) and Lydon et al. (1992, 1993a,b).

In this *Letter* we discuss the first application to the computation of GC isochrones of new results obtained from a grid of detailed 2D radiation-hydrodynamics models including realistic microphysics and a detailed treatment of the radiative transport. A further important feature of these models is that they span a wide range in metallicity ($[M/H] = -2.0$ to $[M/H] = 0.0$; here we adopt the usual spectroscopic notation $[M/H] = \log(M/H)_{\text{star}} - \log(M/H)_{\odot}$, where M is the global metal abundance) and cover the Main Sequence (MS) and lower Red Giant Branch (RGB) region of GC colour-magnitude diagrams.

2. Hydrodynamical models and calibration of the efficiency of convection

The full grid of RHD models is described in detail elsewhere (Ludwig et al. 1998, Freytag et al. 1998a,b), and a comprehensive discussion about the numerical and physical assumptions of the RHD simulations can be found in Ludwig et al. (1994). Here we just recall the main features of the models. Each 2D model describes the atmosphere and upper layers of a star with a convective envelope. It is obtained by solving the time-dependent, non-linear equations of hydrodynamics for a stratified compressible fluid. The calculations take into account a detailed treatment of the equation of state and of the multi-dimensional, non-local, radiative transfer (for more details see Ludwig et al. 1994). Similar to classical model atmosphere calculations, the hydrodynamical models are fully determined by specifying T_{eff} , acceleration of gravity g , and chemical composition, and they lie in the range $4300 \text{ K} \leq T_{\text{eff}} \leq 7100 \text{ K}$, $2.54 \leq \log(g) \leq 4.74$, $-2.0 \leq [M/H] \leq 0.0$.

From this grid of models one can extract the entropy of the deeper, adiabatic convective layers (s_{env}) as a function of T_{eff} , g , and $[M/H]$ (see Ludwig et al. 1998, Freytag et al. 1998a,b). Once this relation is implemented in a stellar evolution code, it completely fixes the T_{eff} of the star as determined from the solution of the stellar structure equations.

A way for implementing easily this dependence of s_{env} on T_{eff} and g into an evolutionary code makes use of the MLT formalism. As explained in detail by Ludwig et al. (1998), for each fixed metallicity one can compute a grid of hydrostatic one-dimensional stellar envelope models based on the MLT, covering the same range of g and T_{eff} spanned by the RHD computations, and using the same input physics. By employing as surface boundary condition the $T(\tau)$ relation

derived from the hydro-models, one can calibrate an effective α (α_{eff}) that is able to reproduce the $s_{\text{env}}-T_{\text{eff}}$ relation obtained from the RHD computations. In this way one can derive a function $\alpha_{\text{eff}}=f(T_{\text{eff}},g)$ at each metallicity, that is easy to use for computing stellar evolutionary models. The estimated error on the values of α_{eff} derived by means of this procedure is equal to ± 0.05 (Ludwig et al. 1998).

Ludwig et al. (1998) discuss the comparison of their RHD model for the solar envelope with the results from helioseismology. They show that the entropy at the bottom of the superadiabatic region as derived from helioseismology would imply a value of α_{eff} for the Sun slightly higher, by $\approx 0.10 \pm 0.05$, than the value deduced from their RHD models. This small discrepancy is explained by comparing their result with the outcome of similar 3D simulations and by the examination of the opacities used in their models. For the Sun the 3D models predict an increase of α by $\approx 0.07 \pm 0.02$ with respect to the 2D ones, and the ATLAS6 opacities (Kurucz 1979) used in the models do not consider the contribution of the molecules. The effect of including this contribution to the opacity would further change α by ≈ 0.1 . The combination of these two effects explains the small discrepancy between the adopted solar RHD model and the results from helioseismology. Another important result derived from the RHD models is that the effect of the envelope He abundance on the derived α_{eff} values is basically negligible.

The calibration of α_{eff} for metal poor stars has been performed by Freytag et al. (1998a,b), and we have used their results for computing isochrones with typical GC metallicities $[M/H]=-2.0$ and $[M/H]=-1.0$ (scaled solar metal distribution), $Y=0.23$, age t ranging from 9 to 14 Gyr, using the code described in Salaris et al. (1997). We have employed the same $T(\tau)$ relation and the same MLT formalism (Böhm-Vitense 1958) used in the calibration of α_{eff} .

We have computed a first set of isochrones using the ATLAS6 low temperature opacities. The only source of possible inconsistencies with the RHD models was in this case the equation of state (EOS) employed in the evolutionary calculations (see Salaris et al. 1997), which is not the same as in the RHD computations. Nevertheless, we have verified that it does not modify appreciably α_{eff} as derived from the RHD calibration. For the sake of comparison, we have computed isochrones for the same age and metallicities, but using a constant, solar calibrated value of α ($\alpha_{\odot}$). Since the ATLAS6 data do not include the contribution of the molecules to the opacity of the stellar matter, we have repeated the same evolutionary computations previously described (with α_{eff} and $\alpha_{\odot}$), using this time the updated Alexander & Ferguson (1994) low temperature opacities, which include the molecular contribution. When using these opacity data, we found a small effect only on the zero point of the $\alpha_{\text{eff}}=f(T_{\text{eff}},g)$ relation. To fix the ideas, the solar calibration with the evolutionary code yields in this case $\alpha=1.69$, while from the RHD models one gets $\alpha=1.59$ for the Sun, that means a deviation by a factor 1.06. This small correction factor (which makes consistent the entropy of the adiabatic layers as derived from RHD models and from helioseismology) for the $\alpha_{\text{eff}}=f(T_{\text{eff}},g)$ relation has therefore been taken into account in the evolutionary calculations.

At this point we have verified that the differences among isochrones computed with α_{eff} and

$\alpha_{\odot}$ are exactly the same in the case of models computed with ATLAS6 or Alexander & Ferguson (1994) opacities. Since the latter data are a more realistic evaluation of the opacity of stellar matter, in the following section we will discuss the isochrones computed using the Alexander & Ferguson (1994) opacities.

Before concluding this section we would like to stress the fact that the derived calibration of α_{eff} is *only intended to reproduce the function $s_{\text{env}}(T_{\text{eff}}, g)$ of the RHD models. The detailed temperature profile and convective velocities of the superadiabatic layers are not represented adequately by the MLT with α_{eff}* , but our main concern here is only the determination of reliable effective temperatures for cool stars.

3. Results and discussion

Representative isochrones (9 and 13 Gyr) for the two considered metallicities, computed with α_{eff} and $\alpha_{\odot}$ are displayed in Figs. 1 and 2, upper panels. The most striking feature is the close resemblance between the two sets of isochrones. The MS loci are coincident, the T_{eff} values of the TO points are very similar (notice the linear scale for the T_{eff} axis), the biggest difference being equal to ≈ 45 K for the 9 Gyr most metal-poor isochrone (an age possibly too young for the metal-poor galactic GC population, see e.g. Salaris & Weiss 1998). Along the RGB the isochrones computed by using α_{eff} are systematically hotter by only ≈ 50 K for $[M/H] = -1.0$ and ≈ 40 K for $[M/H] = -2.0$. To explain this behaviour it is useful to study the run of α_{eff} with respect to $\log(L/L_{\odot})$ along the same isochrones, as shown in the lower panels of the same figures.

The differences between α_{eff} and $\alpha_{\odot}$ along the lower MS are hardly relevant, since the T_{eff} of these stars is insensitive to the choice of α (the entropy jump from the photosphere to the deep adiabatically stratified layers is small anyway), while around the TO they depend on the age of the isochrones. In general, in the youngest, most metal-poor isochrones α_{eff} shows the largest difference with respect to $\alpha_{\odot}$, but since stars in these phases are quite hot ($T_{\text{eff}} \approx 7000$ K) and their convection zones are relatively shallow, the sensitivity of T_{eff} to α is not very large. Along the RGB, where the T_{eff} of stellar models is most sensitive to α because of deeper superadiabatic regions, α_{eff} is systematically higher than $\alpha_{\odot}$ by 0.10-0.15 for both metallicities. This difference causes a systematic shift by ≈ 50 K toward higher T_{eff} with respect to the case of $\alpha_{\odot}$, a quantity marginally significant since the error by ± 0.05 on α_{eff} translates into an error by $\approx \pm 15$ -20 K on the RGB T_{eff}). Qualitatively, the behaviour of isochrones computed using α_{eff} looks similar, for certain features, to the results of the CM formalism; we are referring here to the fact that the TO is cooler (but only for the youngest more metal-poor isochrones) than for the models computed with $\alpha_{\odot}$. But the differences we find are smaller than the predictions of the CM formalism. Moreover, the RGB location in the α_{eff} isochrones is only slightly hotter than in the $\alpha_{\odot}$ ones, while in the case of CM models the RGB is cooler at low metallicity and progressively hotter for increasing metallicities.

At this point, let's turn our attention to the GC age indicators that one can extract from the isochrones. The TO brightness is the most solid one; once the distance is fixed (e.g., from the Horizontal Branch luminosity or by means of the subdwarf fitting technique), the comparison between theoretical and observed TO gives directly the cluster age. The TO colour is also a possible age indicator once the reddening is known, but the present uncertainties on the colour transformations do not favour this method for deriving absolute ages, also if the isochrone T_{eff} and the GC reddenings are determined with high accuracy. A third possibility is to use the reddening- and distance modulus-independent quantity $\Delta(B - V)$, that is the difference in (B-V) between TO and base of the RGB, as defined by Vandenberg et al. (1990). Again, the uncertainties on the colour-transformations prevents from using the absolute value of $\Delta(B - V)$ for deriving absolute GC ages, but the differential use of this quantity is a solid and widely employed indicator of age differences (see, e.g., Vandenberg et al. 1990, Salaris & Weiss 1997), and it is weakly dependent on $[M/H]$.

In Figure 3 (upper and center panel) we compare the TO position (brightness and colour) in the age range 9-14 Gyr for the two sets of isochrones with α_{eff} and $\alpha_{\odot}$. We have transformed the isochrones to the observational V-(B-V) plane according to the colours and bolometric corrections used by Salaris & Weiss (1998), but the results of this comparison do not depend on the particular set of transformations used. As it is evident from the figure, the age differences as derived from the TO brightness (or colour) are basically negligible.

In the lowest panel of Figure 3 we compare $\Delta(B - V)$ as a function of the age, for both sets of isochrones. At each metallicity the two curves corresponding to the two calibrations of the convection lie parallel for all the relevant age range (the absolute values being different by only ≈ 0.02 mag). Therefore, the derivative $\delta(\Delta(B - V))/\delta t$ (and the relative ages derived for the $\Delta(B - V)$) is not affected at all when α_{eff} is used instead of $\alpha_{\odot}$.

In conclusion, the main results of this analysis show that the T_{eff} of GC isochrones computed employing the MLT formalism and $\alpha_{\odot}$, or the α_{eff} calibration as derived from detailed RHD models are in good mutual agreement: the maximum deviations in the relevant age range for galactic GC amounts at most to a systematic shift by $\approx 50 \pm 20$ K along the RGB. As previously discussed, preliminary comparisons (Ludwig et al. 1998) between the adopted grid of 2D RHD models and a small sample of 3D ones show only a very small systematic shift of α_{eff} as derived from the RHD models by ≈ 0.07 , which does not affect our results appreciably.

The next necessary step for finally solving the problem of superadiabatic convection in stellar envelopes involves the computations of 3D model grids with up-to-date equation of state and frequency-dependent opacity tables to improve especially the photospheric temperature structure. Particularly the resolution of the numerical grid in the vertical direction should be improved to resolve the extremely sharp temperature jump at the bottom of the photosphere when the computations are extended to higher luminosities. A better coverage of the transition region from efficient to weak (radiation dominated) convection at high effective temperatures would improve

the base to judge between the MLT and the CM formalism. Nevertheless, the already existing grid of 2D models indicates that in the $T_{\text{eff}}\text{-log } g\text{-[M/H]}$ region of interest for GC stars there is no drastic change in the properties of the envelope convection. Accordingly, the use of the MLT with a constant solar-calibrated $\alpha_{\odot}$ leads only to insignificant errors of at most 0.2 Gyr in the derived ages of globular clusters.

It is important also to remark again that the structure of the superadiabatic convective regions is not suitably reproduced either by $\alpha_{\odot}$ nor by α_{eff} , and that the complete results from RHD models have to be employed whenever a detailed description of the properties of this layers is needed (e.g., for astro- and helioseismology). In addition, the RHD models should be analysed regarding the effects on the T_{eff} -colour-transformations.

We would like to express our gratitude to Hartmut Holweger and Matthias Steffen for continuous support and helpful discussions. We warmly thank John Porter, Achim Weiss, and Hans-Günter Ludwig for reading an early draft of the manuscript and for useful suggestions. B.F. was supported by the *Deutsche Forschungsgemeinschaft*, DFG project Ho 596/36-1 and by the European *Training and Mobility of Researchers* Programme.

REFERENCES

- Alexander, D.R., Ferguson, J.W. 1994, ApJ, 437, 879
- Böhm-Vitense, E. 1958, Zs. Ap., 46, 108
- Canuto, V., Mazzitelli, I. 1991, ApJ, 370, 295
- Chaboyer, B. 1995, ApJ, 444, L9
- Chaboyer, B., Demarque, P., Kernan, P.J., Krauss, L.M. 1998, ApJ, 494, 96
- Deupree, R.G., Varner, T.M. 1980, ApJ, 237, 558
- Freytag, B., Ludwig, H-G., Steffen, M. 1998a, in Workshop on Stellar Structure: Theory and Tests of Convective Energy Transport, ASP series, in press
- Freytag, B., Ludwig, H-G., Steffen, M. 1998b, in preparation
- Frogel, J.A., Persson, S.E., Cohen, J.G. 1981, ApJ, 246, 842
- Gough, D.O., Weiss, N.O. 1976, MNRAS, 176, 589
- Kurucz, R.L. 1979, ApJS, 40, 1
- Ludwig, H-G., Jordan, S., Steffen, M. 1994, A&A, 284, 105
- Ludwig, H-G., Freytag, B., Steffen, M. 1998, A&A, accepted (e-print server astro-ph/9811179)
- Lydon, T.J., Fox, P.A., Sofia, S. 1992, ApJ, 397, 701
- Lydon, T.J., Fox, P.A., Sofia, S. 1993a, ApJ, 403, L79
- Lydon, T.J., Fox, P.A., Sofia, S. 1993b, ApJ, 413, 390
- Mazzitelli, I., D’Antona, F., Caloi, V. 1995, A&A, 302, 382
- Pedersen B.P., Vandenberg D.A., Irwin A.W. 1990, ApJ, 352, 279
- Salaris, M., Weiss, A. 1997, A&A, 327, 107
- Salaris, M., Weiss, A. 1998, A&A, 335, 943
- Salaris, M., Degl’Innocenti, S., Weiss, A. 1997, ApJ, 479, 665
- Vandenberg D.A., Bolte M., Stetson, P.B. 1990, AJ, 100, 445
- Vandenberg D.A., Bolte M., Stetson, P.B. 1996, ARA&A, 34, 461
- Weiss, A., Salaris, M. 1998, A&A, submitted

Fig. 1.— Upper panel: isochrones in the $T_{\text{eff}}\text{-}\log(L/L_{\odot})$ diagram for $[M/H]=-2.0$. Two isochrones ($t=9, 13$ Gyr) computed with α_{eff} are compared to the corresponding isochrones computed with a constant $\alpha_{\odot}$. Lower panel: values of α_{eff} along the hydro-calibrated isochrones; the vertical line represents the constant $\alpha_{\odot}$ of the standard isochrones.

Fig. 2.— As in Figure 1 but for $[M/H]=-1.0$.

Fig. 3.— Upper panel: M_v values for the TO of isochrones computed by using either α_{eff} or $\alpha_{\odot}$, $[M/H]=-2.0$ and -1.0 , $t=9\text{-}14$ Gyr. Central panel: the same, but for the TO (B-V) colour. Lower panel: the same, but for the quantity $\Delta(B - V)$.





